arXiv:1302.2806v1 [quant-ph] 12 Feb 2013

Collapse and Revival and Cat States with an N Spin System

Shane Dooley,[1, *] Francis McCrossan,[1] Derek Harland,[2] Mark J. Everitt,[3] and Timothy P. Spiller[1]
[1] *Quantum Information Science, School of Physics and Astronomy,*
University of Leeds, Leeds LS2 9JT, United Kingdom.
[2] *Department of Mathematical Sciences, Loughborough University,*
Loughborough, Leics LE11 3TU, United Kingdom.
[3] *Department of Physics, Loughborough University,*
Loughborough, Leics LE11 3TU, United Kingdom.

(Dated: February 13, 2013)

We discuss collapse and revival of Rabi oscillations in a system comprising a qubit and a “big spin” (made of N qubits, or spin-1/2 particles). We demonstrate a regime of behaviour analogous to conventional collapse and revival for a qubit-field system, employing spin coherent states for the initial state of the big spin. These dynamics can be used to create a “Schrödinger cat” state of the big spin. Even for relatively small values of N, states with significant potential for quantum metrology applications can result, giving sensitivity approaching the Heisenberg limit.

PACS numbers: 03.65.-w, 42.50.-p, 03.67.-a

Collapse and revival of the Rabi oscillations of a qubit, or two-level atom, coupled to a field mode [1, 2] provide a remarkable and much-discussed illustration of the quantum nature of the composite qubit-field system. The collapse of the Rabi oscillations arises through qubit-field entanglement, yet half way to the revival of the oscillations the field and qubit disentangle again, with the quantum information in their initial states effectively swapped into their counterpart system [3]. This enables, for example, the generation of so-called “Schrödinger cat” states of a field mode—superpositions of two (approximate) coherent states of the field [4]. The addition of further qubits has led to the phenomena of entanglement sudden death [5–7], collapse and revival of entanglement [7] and cat-swapping [8].

In this work we discuss collapse and revival of the Rabi oscillations of a qubit coupled to a “big spin” made of N qubits, or spin-1/2 particles. We demonstrate that preparation of the N spins in a spin coherent state—the analogue of a coherent state for a field mode—facilitates collapse and revival phenomena. We isolate a regime of parameter space where conventional looking collapse and revival occurs and thus “Schrödinger cat” states of the big spin emerge prior to revival, although we note that different and interesting forms of revival dynamics occur in other parameter regimes.

Quantum metrology [9–11] is an emerging quantum information application where entangled quantum resources are employed to measure an unknown external potential or field that through interaction generates a phase in the state of the quantum resources. Enhanced measurement precision—beyond the “Standard Quantum Limit” achieved using the resources one-by-one, or classically—is possible. In principle, precision all the way down to the “Heisenberg limit” can be attained. Schrödinger cat states are one form of non-classical resource with significant potential for metrology, with spin cats enabling enhanced magnetic field sensing [12]. We therefore assess the ability of the states produced through spin collapse and revival for quantum metrology, and demonstrate that accuracy close to that of the Heisenberg limit could be attained.

We begin our discussion with the familiar qubit-field dynamics. The Jaynes-Cummings (JC) model [1] for the interaction of a bosonic field mode with a qubit, or two level atom, is described by the Hamiltonian

$$H_{\mathrm{JC}} = \omega \hat{a}^\dagger \hat{a} + \frac{\Omega}{2}\hat{\sigma}_z + \lambda\left(\hat{a}^\dagger \hat{\sigma}_- + \hat{a}\hat{\sigma}_+\right), \qquad (1)$$

where $\hat{a}^\dagger$ and $\hat{a}$ are the field mode creation and annihilation operators respectively, and $\hat{\sigma}_z = |0\rangle\langle 0| - |1\rangle\langle 1|$, $\hat{\sigma}_+ = |0\rangle\langle 1|$ and $\hat{\sigma}_- = |1\rangle\langle 0|$ are qubit operators. Assuming resonance ($\Omega = \omega$), if the qubit is initially in the state $|0\rangle$ and the field initially in a coherent state $|\alpha\rangle = \left||\alpha|e^{i\phi}\right\rangle$ we see the well-known collapse and revival of oscillations of $\langle\hat{\sigma}_z(t)\rangle$ [red line, figure 1 (a)] and the dip in the qubit entropy at half of the revival time (green line, figure 1 (a)).

Gea-Banacloche [3, 4] has made an insightful analysis of these features. For example, he has shown that when $|\alpha|^2 \gg 1$, there are two initially orthogonal qubit states, $|D_\pm(0)\rangle$, that – to a good approximation – evolve without entangling with the field mode. He has also shown that at a particular time t_0, which is equal to half the revival time, these states coincide: $|D_+(t_0)\rangle = |D_-(t_0)\rangle$. This is known as the *attractor state* of the qubit and it is independent of the initial qubit state. Since any pure state of the qubit can be written as a superposition of $|D_+(0)\rangle$ and $|D_-(0)\rangle$, it follows that any initial qubit state will converge to the attractor state at t_0. This explains the dip in qubit entropy at half the revival time in figure 1 (a). Since the state of the composite qubit-field system is pure at all times, the field must also be in a pure state at t_0. For a judicious choice of initial qubit state, the field is in a “Schrödinger cat” state at this time [4, 13].

Instead of the interaction of a field mode with a two level atom, we consider the interaction of a system of N spin-$\frac{1}{2}$ particles (which we refer to as "the big spin") with a single spin-$\frac{1}{2}$ particle (a qubit) via the Hamiltonian

$$H = \omega J_z + \frac{\Omega}{2}\sigma_z + \frac{\lambda}{\sqrt{N}}\left(J_+\sigma_- + J_-\sigma_+\right), \tag{2}$$

where $J_z \equiv \frac{1}{2}\sum_{i=1}^{N}\sigma_z^{(i)}$ and $J_\pm \equiv \sum_{i=1}^{N}\sigma_\pm^{(i)}$ are operators that act on the big spin and $\sigma_z^{(i)} = \left|\uparrow^{(i)}\right\rangle\left\langle\uparrow^{(i)}\right| - \left|\downarrow^{(i)}\right\rangle\left\langle\downarrow^{(i)}\right|$ acts on the individual spins that make up the big spin.

We say that the big spin is initially in a *spin coherent state* $|N,\zeta\rangle$. A spin coherent state [14, 15] of the N spin system is one in which each of the N spins is in the same pure state. This state is parameterised by the complex number ζ:

$$|N,\zeta\rangle = \bigotimes_{i=1}^{N}\left(\frac{1}{\sqrt{1+|\zeta|^2/N}}\left|\downarrow^{(i)}\right\rangle + \frac{\zeta/\sqrt{N}}{\sqrt{1+|\zeta|^2/N}}\left|\uparrow^{(i)}\right\rangle\right). \tag{3}$$

There is an equivalent definition of the spin coherent state in terms of the angular momentum states, $\left|\frac{N}{2}, n-\frac{N}{2}\right\rangle \equiv |n\rangle$, which are simultaneous eigenstates of $\hat{J}_z$ (with eigenvalue $n - N/2$) and the total angular momentum operator $\hat{J}^2 = \hat{J}_x^2 + \hat{J}_y^2 + \hat{J}_z^2$ [with eigenvalue $\frac{N}{2}(\frac{N}{2}+1)$]. In terms of these states, the spin coherent state is $|N,\zeta\rangle = \sum_{n=0}^{N} C_n|n\rangle$ where

$$C_n = \frac{1}{\left(1+\frac{|\zeta|^2}{N}\right)^{N/2}}\sqrt{\frac{N!}{(N-n)!n!}}\left(\frac{\zeta}{\sqrt{N}}\right)^n. \tag{4}$$

Plotted in figure 1 (b) (again assuming resonance $\Omega = \omega$, qubit initially in $|0\rangle$, and initial spin coherent state with $\zeta = 4$, $N = 170$) are $\langle\sigma_z(t)\rangle$ and the linear entropy of the qubit. The oscillations in $\langle\sigma_z\rangle$ undergo a collapse and revival that is very similar to the collapse and revival in the JC model.

The similarities between our qubit-big spin system and the JC model can be understood by looking at the $N \to \infty$ limit of the big spin system. To see the connection, we write $\hat{J}_\pm$ in terms of the angular momentum states:

$$\frac{\hat{J}_+}{\sqrt{N}} = \sum_{n=0}^{N}\sqrt{(n+1)\left(1-\frac{n}{N}\right)}\,|n+1\rangle\langle n|\ ; \tag{5}$$

$$\frac{\hat{J}_-}{\sqrt{N}} = \sum_{n=0}^{N}\sqrt{n\left(1-\frac{n-1}{N}\right)}\,|n-1\rangle\langle n|\,. \tag{6}$$

Taking the $N \to \infty$ limit we find that $\frac{\hat{J}_+}{\sqrt{N}}$ and $\frac{\hat{J}_-}{\sqrt{N}}$ are mathematically identical to the bosonic creation and annihilation operators with the angular momentum states $|n\rangle \equiv \left|\frac{N}{2}, n-\frac{N}{2}\right\rangle$ playing the role of "Fock states" in our big spin system. Similarly, it can be shown that $\lim_{N\to\infty}\left(\hat{J}_z + \frac{N}{2}\right)$ is mathematically the same as $\hat{a}^\dagger\hat{a}$, the field mode number operator. Combining these facts we see that our Hamiltonian (2) is equivalent to the Jaynes-Cummings Hamiltonian (1) in the $N \to \infty$ limit. Moreover, given our definition (4) of the spin coherent state, one can use the Poisson Limit Theorem [16] to show that, in the $N \to \infty$ limit (if $|\zeta|^2$ is kept constant as we take the limit), the spin coherent state is mathematically identical to the field mode coherent state:

$$\lim_{N\to\infty}|N,\zeta\rangle = e^{-|\zeta|^2/2}\sum_{n=0}^{\infty}\frac{\zeta^n}{\sqrt{n!}}|n\rangle\,. \tag{7}$$

In the $N \to \infty$ limit we see collapse and revival of $\langle\sigma_z\rangle$ because both our big spin initial state and Hamiltonian are mathematically the same as those that result in collapse and revival in the field-qubit interaction. This correspondence between our big spin-qubit model and the JC model is exact in the $N \to \infty$ limit. If N is finite, it is not exact but is very good when $\frac{|\zeta|^2}{N} \ll 1 \ll N$ (assuming that the big spin is initially in the spin coherent state $|N,\zeta\rangle$). If $|\zeta|^2 \approx N$ the correspondence between the qubit-big spin model and the JC model breaks down.



FIG. 1. (colour online). When $\frac{|\zeta|^2}{N} \ll 1 \ll N$, there is a correspondence between the qubit-big spin model and the JC model. Here we see collapse and revival of $\langle\sigma_z\rangle$ (red line) in (a) the JC model, and (b) the qubit-big spin model.

We make use of the correspondence when $\frac{|\zeta|^2}{N} \ll 1 \ll N$ to propose a method of creating spin cat states. By analogy with the approximation of Gea-Banacloche for the JC model, we suggest that initial qubit states $|D_\pm(0)\rangle = \frac{1}{\sqrt{2}}\left(|0\rangle \pm e^{-i\phi}|1\rangle\right)$ evolve without entangling with the big spin when $1 \ll |\zeta|^2 \ll N$:

$$|D_\pm(0)\rangle|N,\zeta\rangle \to |D_\pm(t)\rangle\, e^{\mp i\lambda t\sqrt{\hat{J}_-\hat{J}_+/N}}|N,\zeta\rangle\,, \tag{8}$$

where $|D_\pm(t)\rangle = \frac{1}{\sqrt{2}}\left(|0\rangle \pm e^{-i\phi}e^{\pm i\pi t/2t_0}|1\rangle\right)$. Here $t_0 \equiv \frac{\pi|\zeta|}{\lambda}\sqrt{1-\frac{|\zeta|^2}{N}}$ and ϕ is the complex phase of $\zeta = |\zeta|e^{i\phi}$.

At time $t = t_0$ we can see that $|D_+(t_0)\rangle = |D_-(t_0)\rangle$ and the qubit is in the attractor state $\frac{1}{\sqrt{2}}\left(|0\rangle + ie^{-i\phi}|1\rangle\right)$, regardless of the initial state of the qubit. The big spin is, at this time, in a pure state that will depend on the initial state of the qubit. For initial qubit state $|0\rangle = \frac{1}{\sqrt{2}}\left(|D_+(0)\rangle + |D_-(0)\rangle\right)$, the big spin is, at time t_0, in the state

$$|\psi_\zeta\rangle = \frac{1}{\sqrt{2\mathcal{M}}}\left[e^{-i\lambda t_0\sqrt{\frac{\hat{J}_-\hat{J}_+}{N}}}|N,\zeta\rangle + e^{+i\lambda t_0\sqrt{\frac{\hat{J}_-\hat{J}_+}{N}}}|N,\zeta\rangle\right], \tag{9}$$

a "Schrödinger cat" state. The $\mathcal{M}$ has been introduced to maintain normalisation of $|\psi_\zeta\rangle$ since $e^{-i\lambda t_0\sqrt{\hat{J}_-\hat{J}_+/N}}|N,\zeta\rangle$ and $e^{+i\lambda t_0\sqrt{\hat{J}_-\hat{J}_+/N}}|N,\zeta\rangle$ are, in general, not orthogonal to each other.



FIG. 2. (colour online). (a) Fidelity at t_0 is marked by a black dot. For $N = 40, 70, 100$ the fidelity is high at this time. The fidelity around t_0 is highly oscillatory. ($|\zeta|^2 = 6.5$ in each case.); (b) For N not too small, fidelity at t_0 is high for a broad range of values of $|\zeta|^2 \ll N$. Our approximation is no longer valid in the region where $|\zeta|^2 \approx N$.

This approximation is found by analogy with Gea-Banacloche's JC model analysis. To rigorously show that it is a good approximation, one could show that the fidelity of the approximation against the exact state approaches unity as $|\zeta|^2$ and N both increase (with $1 \ll |\zeta|^2 \ll N$). Since this is a lengthy calculation we support our claim here by pointing to the fidelity plots, figures 2(a) and 2(b). These plots show $F = \sqrt{\langle\psi_\zeta|\rho_{\text{BS}}(t)|\psi_\zeta\rangle}$, the fidelity of $|\psi_\zeta\rangle$ against the (exact) reduced big spin state $\rho_{\text{BS}}(t)$ plotted against time [figure 2(a)], and against $|\zeta|^2$ at t_0 [figure 2(b)], for $N = 10, 40, 70, 100$. For $N = 100$, for example, the fidelity at t_0 [marked by a black dot in figure 2(a)] is high (~ 0.95). This is telling us that the big spin system is close to the cat state $|\psi_\zeta\rangle$ at t_0. Figure 2(b) shows how the fidelity at t_0 varies with $|\zeta|^2$. For $N = 100$, there is a broad range of values of $|\zeta|^2 \ll N$ for which the fidelity at t_0 is high and slowly varying. This tells us that the fidelity at t_0 is not too sensitive to our choice of $|\zeta|^2$ for the initial spin coherent state. The fast oscillation of the fidelity in figure 2(a), however, tells us that the fidelity is sensitive to the interaction time.

Although we expect to see a good cat state when $1 \ll |\zeta|^2 \ll N$ (because of the correspondence between our big spin model and the JC model), numerics show that, interestingly, fidelity to a cat state is high for relatively small values of N. At $N = 40$, for example (green lines in figure 2), $F \geq 0.9$ at t_0, again for a broad range of values of $|\zeta|^2 \ll N$.

To see that this is indeed a cat state, we plot in figure 3, for $N = 5, 10, 20, 40$, the spin Wigner functions [17] of $\rho_{\text{BS}}(t_0)$. A spin coherent state is represented by a circular blob in a spin Wigner plot. A superposition of spin coherent states would be represented by two circular blobs with interference fringes between them. Here, in figure 3(d), instead of circles, we have two crescent shapes with interference fringes between them – clearly a cat state, although not quite a superposition of spin coherent states.

We now try to quantify the usefulness of such a state for metrology or, in particular, for magnetic field sensing. A system of N spin-$\frac{1}{2}$ particles initially in the cat state $|\psi_\zeta\rangle$ is allowed to interact with an unknown (classical) magnetic field, $\vec{B} = B_y$, via the Hamiltonian $\hat{H} = \gamma B_y \hat{J}_y$, where γ is the gyromagnetic ratio of our N spin system. After a time t, our spin system is in the state $|\psi_\zeta(\theta)\rangle = e^{i\gamma B_y t \hat{J}_y}|\psi_\zeta\rangle = e^{i\theta\hat{J}_y}|\psi_\zeta\rangle$, where we have defined $\theta = \gamma t B_y$. Since γ and t are assumed to be known, estimating θ is the same as estimating B_y. The precision $\Delta\theta$ with which we can estimate the parameter θ is bounded by the Cramer-Rao inequality [19], $(\Delta\theta)^2 \geq 1/\mathcal{F}$, where $\mathcal{F}$ is the quantum Fisher information. For ease of comparison between different values of N we quantify precision by $N(\Delta\theta)^2 \geq N/\mathcal{F}$. Given that our N spin system evolves unitarily and is initially in pure state $|\psi_\zeta\rangle$ we can write the quantum Fisher information as

$$\mathcal{F} = 4\left(\Delta\hat{J}_y\right)^2 = \langle\psi_\zeta|\hat{J}_y^2|\psi_\zeta\rangle - \langle\psi_\zeta|\hat{J}_y|\psi_\zeta\rangle^2. \tag{10}$$

In figure 4 we plot $N/\mathcal{F}$ against $|\zeta|^2/N$ for $N = 10, 40, 70, 100$. If $\zeta = 0$, our initial state $|\psi_{\zeta=0}\rangle$ is just a spin coherent state and $N/F = 1$, the standard quan-


(a) $N = 5$ (b) $N = 10$

(c) $N = 20$ (d) $N = 40$

FIG. 3. (colour online). Spin Wigner functions of $\rho_{\mathrm{BS}}(t_0)$, the exact reduced big spin state at t_0 (with $|\zeta|^2 = 0.16N$). Interactive figures showing the states in (b) and (c) are available as supplementary material [18].



FIG. 4. (colour online). $N/\mathcal{F} = 1$ at $|\zeta|^2 = 0$ corresponding to the standard quantum limit. The Heisenberg limit, $N/\mathcal{F} = 1/N$ is marked by a black line.

tum limit. The Heisenberg limit, $N/\mathcal{F} = 1/N$, is marked in figure 4 by a black line for each N . We see that, especially for large N, our cat state $|\psi_\zeta\rangle$ can allow for magnetic field sensing close to the Heisenberg limit, even in the $|\zeta|^2 \ll N$ regime in which the cat state emerges from the collapse and revival dynamics.

In conclusion, we have considered collapse and revival phenomena for a single spin coupled to a composite big spin, identifying a parameter regime corresponding to conventional qubit-field mode behaviour. Here the evolving system can produce non-classical (Schrödinger cat type) states of the big spin. Such states are capable of quantum-enhanced field sensing that approaches the Heisenberg limit for modest values of N, the number of component spins in the big spin. From a practical perspective, these states can be generated from very straightforward initial states: the qubit is initially in a pure state and the big spin is in a separable state of its N component spins with each of the spins aligned (a spin coherent state). Furthermore, any physical realisation for which there is control at the Rabi period time scale should have sufficient time resolution to identify the time(s) at which a cat is generated.

Future work will consider in detail the robustness of the collapse and revival and cat state generation to various forms of decoherence. In addition, in the $|\zeta|^2 \approx N$ parameter regime it is possible to explore other regimes of “non-standard” collapse and revival. These will be discussed in a forthcoming paper [20].

---

* pysd@leeds.ac.uk

[1] E. Jaynes and F. Cummings, Proc. IEEE. **51** (1963).
[2] J. H. Eberly, N. B. Narozhny, and J. J. Sanchez-Mondragon, Phys. Rev. Lett. **44**, 1323 (1980).
[3] J. Gea-Banacloche, Phys. Rev. Lett. **65**, 3385 (1990).
[4] J. Gea-Banacloche, Phys. Rev. A **44**, 5913 (1991).
[5] T. Yu and J. H. Eberly, Phys. Rev. Lett. **93**, 140404 (2004).
[6] T. Yu and J. H. Eberly, Science **323**, 598 (2009).
[7] C. Jarvis, D. Rodrigues, B. Gyorffy, T. Spiller, A. Short, and J. Annett, New J. Phys. **11**, 103047 (2009).
[8] C. E. A. Jarvis, D. A. Rodrigues, B. L. Györffy, T. P. Spiller, A. J. Short, and J. F. Annett, J. Opt. Soc. Am. B **27**, A164 (2010).
[9] H. Lee, P. Kok, and J. P. Dowling, Journal of Modern Optics **49**, 2325 (2002).
[10] V. Giovannetti, S. Lloyd, and L. Maccone, Science **306**, 1330 (2004).
[11] V. Giovannetti, S. Lloyd, and L. Maccone, Nature Photonics **5**, 222 (2011).
[12] H.-N. Xiong, J. Ma, W.-F. Liu, and X. Wang, Quantum Info. Comput. **10**, 498 (2010).
[13] V. Bužek, H. Moya-Cessa, P. Knight, and S. Phoenix, Physical Review A **45**, 8190 (1992).
[14] F. Arecchi, E. Courtens, R. Gilmore, and H. Thomas, Physical Review A **6**, 2211 (1972).
[15] J. M. Radcliffe, Journal of Physics A: General Physics **4**, 313 (1971).
[16] A. Papoulis and S. Pillai, *Probability, random variables, and stochastic processes*, McGraw-Hill electrical and electronic engineering series (McGraw-Hill, 2002).
[17] D. Harland, M. Everitt, K. Nemoto, T. Tilma, and T. Spiller, Phys. Rev. A **86**, 062117 (2012).
[18] See Supplemental Material at [URL] for interactive figures illustrating the same states as figures 3(b) (N=10) and 3(c) (N=20). Plotted are spherical spin Wigner functions. See [17] for more details.
[19] S. L. Braunstein and C. M. Caves, Phys. Rev. Lett. **72**, 3439 (1994).
[20] S. Dooley, F. McCrossan, D. Harland, M. Everitt, and T. Spiller, *In preparation* (2013).